A novel KdV-type equation ?

Sandra CARILLO
Dipartimento Scienze di Base e Applicate per l'Ingegneria
Sapienza - Università di Roma, 16, Via A. Scarpa, 00161 Rome, Italy
&
National Institute for Nuclear Physics (INFN), Rome, Italy
Gr. Roma1, IV - Mathematical Methods in NonLinear Physics
Thursday 23rd February, 2017

arXiv:1702.06874v1 [math.AP] 22 Feb 2017

Abstract

A *novel* third order nonlinear evolution equation is introduced. It is connected, via Bäcklund transformations, with the Korteweg-deVries (KdV), modified Korteweg-deVries (mKdV) equation and other third order nonlinear evolution equations. Hence, it is termed KdV-type equation. In particular, the Bäcklund transformation which links the KdV singularity manifold equation with the introduced equation is used to show that the latter, further to enjoy a nontrivial invariance property, admits a hereditary recursion operator. Hence, results are extended to the generated whole hierarchy of nonlinear evolution equations. The established links allow to extend the Bäcklund chart in [22] to include also the newly introduced equation, which, then, turns out to be related also to the Dym equation.

Keywords: Nonlinear Evolution Equations; Bäcklund Transformations; Recursion Operators; Korteweg deVries-type equations; Cole-Hopf Transformations.

AMS Classification: 58G37; 35Q53; 58F07

1 Introduction

The relevance of Bäcklund transformations in soliton theory is well established, see [2, 42, 39, 41, 27] where a wide variety of applications of Bäcklund and Darboux Transformations and their connections with partial differential equations admitting soliton solutions is given. Here, the concern is on Bäcklund transformations as a tool to investigate structural properties of nonlinear evolution equations. Specifically, the Bäcklund chart in [22] is reconsidered to incorporate a further[1] third order nonlinear equation, termed KdV-type since is connected via Bäcklund transformations with the Korteweg deVries (KdV), the modified Korteweg deVries (mKdV), the *Korteweg deVries interacting soliton* (int.sol.KdV)[20] and the *Korteweg deVries singularity manifold*

[1]The author believes in its novelty since it is not comprised in the examined literature which includes classification works such as [1, 48, 33, 34].

(KdV sing.)[2] equations. The introduced equation is proved to enjoy an invariance property. In addition, since connected via Bäcklund transformations to the other KdV-type equations, according to [18, 19], admits a hereditary recursion operator. The heritariness of all the recursion operators admitted by the equations in the Bäcklund chart allow to extend all the links to the whole corresponding hierarchies; hence, previous results [22] are generalized. Further subsequent generalizations of links in [22] to non-Abelian equations are comprised in [7, 8, 11, 13].

The material is organized as follows. The opening Section 2 is devoted to recall the definition of Bäcklund transformation adopted throughout this work together with its consequences which are most relevant to the present investigation. In the following Section 3 the *novel* nonlinear evolution equation is introduced. Specifically, it is shown to be linked, via suitable Bäcklund transformations with the mKdV and the KdV singularity manifold equations. Notably, both the equations introduced in [13] reduce to this equation when commutativity is assumed. In Section 4, and 5, respectively, the introduced KdV-type equation is proved to enjoy an invariance property and is inserted in the Bäcklund chart in [22], which is, then, further extended. In Section 6, via the links in the Bäcklund chart, the studied equation is shown to admit a hereditary recursion operator. The latter is constructed and, hence the corresponding hierarchy follows. As a consequence [18], the generated whole hierarchy of nonlinear evolution equations is connected to all the hierarchies in the Bäcklund chart. Concluding remark as well as how the present work is related to previous ones and, in particular, with the research progam devoted to the study of non-Abelian nonlinear evolution equations [7, 8, 11, 13], are comprised in the closing Section 7.

2 Some background definitions

This section is devoted to provide some background definitions which are of use throughout the whole article. Since many definitions are not unique in the literature, here, those ones here adopted are provided.

First of all, the notion of Bäcklund Transformation, according to Fokas and Fuchssteiner [18] is recalled (see also the book by Rogers and Shadwick [42]). Consider non linear evolution equations of the type

$$u_t = K(u) \tag{2.1}$$

where the unknown function u depends on the independent variables x, t and, for fixed t, $u(x,t) \in M$, a manifold modeled on a linear topological space so

[2]The Korteweg deVries singuarity manifold equation, introduced in [49], is also known as as UrKdV or Schwarz-KdV [17, 50].

that the generic *fiber* T_uM, at $u \in M$, can be identified with M itself [3]. and $K : M \to TM$, is an appropriate C^∞ vector field on a manifold M to its tangent manifold TM. Let, now

$$v_t = G(v) \tag{2.2}$$

denote another nonlinear evolution equation. If it assumed $u(x,t) \in M_1$ and $v(x,t) \in M_2$ where M_1, M_2 represent manifolds modeled on a linear topological space so that, $K : M_1 \to TM_1$ and $G : M_2 \to TM_2$ represent appropriate C^∞-vector fields on the manifolds $M_i, i = 1, 2$, then

$$u_t = K(u), \quad K : M_1 \to TM_1, \quad u : (x,t) \in \mathbb{R} \times \mathbb{R} \to u(x,t) \in M_1 \tag{2.3}$$
$$v_t = G(v), \quad G : M_2 \to TM_2, \quad v : (x,t) \in \mathbb{R} \times \mathbb{R} \to v(x,t) \in M_2. \tag{2.4}$$

Here, according to the usual choice when soliton solutions are considered, it is further assumed $M := M_1 \equiv M_2$. Then, [18] a Bäcklund transformation can defined as follows.

Definition *Given two evolution equations, $u_t = K(u)$ and $v_t = G(v)$, then $B\ (u\ ,\ v) = 0$ represents a Bäcklund transformation between them whenever given two solutions of such equations, say, respectively, $u(x,t)$ and $v(x,t)$ such that*

$$B(u(x,t), v(x,t))|_{t=0} = 0 \tag{2.5}$$

it follows that,

$$B(u(x,t), v(x,t))|_{t=\bar{t}} = 0, \quad \forall \bar{t} > 0\ , \quad \forall x \in \mathbb{R}. \tag{2.6}$$

Hence, solutions admitted by the two equations are connected via the Bäcklund transformation which establishes a correspondence between them: it can graphicallly represented as

$$\boxed{u_t = K(u)} \overset{B}{—} \boxed{v_t = G(v)}\ . \tag{2.7}$$

In addition, if, the nonlinear evolution equation (2.1) admits a hereditary recursion operator [18, 38], , denoted as $\Phi(u)$, it can be written as

$$u_t = \Phi(u) u_x \quad \text{where} \quad K(u) = \Phi(u) u_x\ . \tag{2.8}$$

The Bäcklund transformation allows to construct the operator $\Psi(v)$

$$\Psi(v) = \Pi\, \Phi(u)\, \Pi^{-1} \tag{2.9}$$

where

$$\Pi := -B_v^{-1} B_u\ \ ,\ \ \Pi : TM_1 \to TM_2\ , \tag{2.10}$$

[3] It is generally assumed that M is the space of functions $u(x,t)$ which, for each fixed t, belong to the Schwartz space S of *rapidly decreasing functions* on $\mathbb{R}^n$, i.e. $S(\mathbb{R}^n) := \{f \in C^\infty(\mathbb{R}^n) : \|f\|_{\alpha,\beta} < \infty, \forall \alpha, \beta \in \mathbb{N}_0^n\}$, where $\|f\|_{\alpha,\beta} := sup_{x \in \mathbb{R}^n} \left|x^\alpha D^\beta f(x)\right|$, and $D^\beta := \partial^\beta / \partial x^\beta$; throughout this article $n = 1$.

while B_u and B_v denote the Frechet derivatives of the Bäcklund transformation $B(u,v)$. Then, [18] , the operator $\Psi(v)$ represents the hereditary recursion operator admitted by the equation $v_t = G(v)$ which, thus, can be written under the form

$$G(v) = \Psi(v)\, v_x \ .$$

That is, according to [18], given the Bäcklund transformation $B(u,v)$, and the hereditary recursion operator $\Phi(u)$ admitted by equation (2.1), then, also equation (2.2) admits a hereditary recursion operator: it is obtained on use of the operator Π, (2.10), via the trasformation formula (2.9).

On subsequent applications of the admitted recursion operators are, respectively, the two hierarchies

$$u_t = [\Phi(u)]^n\, u_x \qquad \text{and} \qquad v_t = [\Psi(v)]^n\, v_x \ , \quad n \in \mathbb{N} \tag{2.11}$$

of evolution equations can be constructed [19]; their *base members* equations, which correspond to $n = 1$, coincide with equations (2.1) and (2.2). Fixed any $n_0 \in \mathbb{N}$, the two equations $u_t = [\Phi(u)]^{n_0}\, u_x$ and $v_t = [\Psi(v)]^{n_0}$ are connected, via the same Bäcklund Transformation which connects the two base members equations. This extension to the whole hierarchies is graphically represented by the following Bäcklund chart

$$\boxed{u = [\Phi(u)]^n\, u_x} \overset{B}{---} \boxed{v_t \ = \ [\Psi(v)]^n\, v_x} \ . \tag{2.12}$$

which emphasizes that the link between the two equations (2.1) and (2.2) is extended to corresponding members of the two hierarchies generated, respectively, by the recursion operators Φ and Ψ.

3 A third order KdV-type equation

In this Section a new third order nonlinear evolution equation is presented. Then, it is proved to be connected with all the equations in the Bäcklund chart constructed in [22]. Consider the nonlinear evolution equation:

$$w_t = w_{xxx} - 3\frac{w_x w_{xx}}{w} \ . \tag{3.1}$$

The following proposition can be proved.

Proposition 1
Equation (3.1) is linked to the mKdV equation

$$v_t = v_{xxx} - 6v^2 v_x \tag{3.2}$$

via the Cole-Hopf [16, 28] *transformation*

$$\text{CH}: \qquad vw - w_x = 0 \ . \tag{3.3}$$

Proof
On substitution, in (3.2), of v in terms of w according to the latter gives:

$$\left(\frac{w_x}{w}\right)_t = \left(\frac{w_x}{w}\right)_{xxx} - 6\left(\frac{w_x}{w}\right)^2\left(\frac{w_x}{w}\right)_x$$

which, since the assumed regularity of v and w implies Schwartz theorem on order of partial derivation holds, delivers

$$\left(\frac{w_t}{w}\right)_x = \left[\frac{w_{xxx}}{w} - 3\frac{w_x w_{xx}}{w^2} + 2\left(\frac{w_x}{w}\right)^3 - 2\left(\frac{w_x}{w}\right)^3\right]_x$$

hence, after simplification, equation (3.1) follows. □

Proposition 2
Equation (3.1) is linked to the KdV sing manifold equation

$$\varphi_t = \varphi_x\{\varphi; x\} \quad , \quad \{\varphi; x\} = \left(\frac{\varphi_{xx}}{\varphi_x}\right)_x - \frac{1}{2}\left(\frac{\varphi_{xx}}{\varphi_x}\right)^2 \tag{3.4}$$

via the Bäcklund transfornation

$$\mathrm{B}: \qquad w^2 - \varphi_x = 0 \ . \tag{3.5}$$

Proof
The Bäcklund transformation (3.5) implies:

$$\begin{aligned} 2ww_t &= \varphi_{xt} \\ 2\frac{w_{xx}}{w} &= \frac{\varphi_{xxx}}{\varphi_x} - \frac{1}{2}\frac{\varphi_{xx}^2}{\varphi_x^2} \\ 2\frac{w_{xxx}}{w} &= \frac{\varphi_{xxxx}}{\varphi_x} - \frac{3}{2}\frac{\varphi_{xx}\varphi_{xxx}}{\varphi_x^2} + \frac{3}{4}\frac{\varphi_{xx}^3}{\varphi_x^3} \end{aligned}$$

Substitution of the latter in (3.1) gives:

$$\varphi_{xt} = \varphi_{xxxx} - 3\frac{\varphi_{xx}\varphi_{xxx}}{\varphi_x} + \frac{3}{2}\frac{\varphi_{xx}^3}{\varphi_x^2}$$

since

$$\{\varphi; x\}_x = \varphi_{xxxx} - 3\frac{\varphi_{xx}\varphi_{xxx}}{\varphi_x} + \frac{3}{2}\frac{\varphi_{xx}^3}{\varphi_x^2}$$

on integration with respect to x, the KdV sing manifold equation (3.4) follows and the proof is complete. □

Remark
Both the two new non-Abelian nonlinear evolution equations[4]

$$W_t = W_{xxx} - 3\,W_{xx}\,W^{-1}W_x \ , \quad Z_t = Z_{xxx} - 3\,Z_x\,Z^{-1}Z_{xx} \tag{3.6}$$

introduced in [13], reduce to equation (3.1) when the assumption of a non-Abelian unknown is removed.

[4]Capital case is used to emphasize that the unknown fuctions Z and W are non-Abelian ones, in [13] operators on a Banach space.

4 A non trivial invariance

Some properties the introduced equation (3.1) enjoys are studied in this section and in the following ones. In particular, this section is devoted to prove an invariance property it enjoys.

First of all, equation (3.1) is *scaling invariant* since substitution of $\alpha w, \forall \alpha \in \mathbb{C}$, to w leaves it unchanged. In addition, the following the following proposition holds.

Proposition 2
The nonlinear evolution equation (3.1) *is invariant under the transformation*

$$\mathrm{I}: \quad \hat{w}^2 = \frac{ad-bc}{(D^{-1}(w^2)+d)^2} w^2, \quad a,b,c,d \in \mathbb{C} \ s.t. \ ad-bc \neq 0, \tag{4.1}$$

where

$$D^{-1} := \int_{-\infty}^{x} d\xi$$

is well defined since so called *soliton solutions* are looked for in the Schwartz space $S(\mathbb{R}^n)$ [5].

Proof
The KdV singularity manifold equation (3.4) is invariant under the Möbius group of transformations

$$\mathrm{M}: \quad \hat{\varphi} = \frac{a\varphi + b}{c\varphi + d}, \qquad a,b,c,d \in \mathbb{C} \qquad \text{such that} \quad ad-bc \neq 0. \tag{4.2}$$

Recalling that, according to proposition 2, the KdV singularity manifold equation (3.4) and equation (3.1) are connected to each other via the Bäcklund transformation B (3.5), the result is readily obtained. Indeed, the following Bäcklund chart

$$\begin{array}{lcc}
\mathrm{M}: \hat{\varphi} = \dfrac{a\varphi+b}{c\varphi+d} & \boxed{w_t = w_{xxx} - 3\dfrac{w_x w_{xx}}{w}} & \xrightarrow{w^2-\varphi_x=0} \quad \boxed{\varphi_t \ = \ \varphi_x\{\varphi; x\}} \\
& \updownarrow \ \mathrm{I} & \updownarrow \ \mathrm{M} \\
\forall a,b,c,d \in \mathbb{C} |\ ad-bc \neq 0 & \boxed{\hat{w}_t = \hat{w}_{xxx} - 3\dfrac{\hat{w}_x \hat{w}_{xx}}{\hat{w}}} & \xrightarrow{\hat{w}^2-\hat{\varphi}_x=0} \quad \boxed{\hat{\varphi}_t \ = \ \hat{\varphi}_x\{\hat{\varphi}; x\}}
\end{array}$$

shows that the invariance I is obtained via composition of the Möbius transformation M with the two Bäcklund transformations

$$\mathrm{B}: \quad w^2 - \varphi_x = 0 \quad \text{and} \qquad \widehat{\mathrm{B}}: \quad \hat{w}^2 - \hat{\varphi}_x = 0 \ . \qquad \square$$

[5] see footnote on page 4.

5 Extended Bäcklund chart

In this section, the equation (3.1) is inserted in the Bäcklund chart in [22] which, then, is further extended. Indeed, combination of the two transformations CH and B in the previos section gives

$$v - \frac{1}{2}\frac{\varphi_{xx}}{\varphi_x} = 0 \tag{5.1}$$

which coincides with the link in [22] between mKdV and KdV sing. equations[6]. The links given so far are sumarized in the following Bäcklund chart:

$$\boxed{\text{KdV}(u)} \overset{(a)}{—} \boxed{\text{mKdV}(v)} \overset{(b)}{—} \boxed{\text{new eq}(w)} \overset{(c)}{—} \boxed{\text{KdV sing.}(\varphi)} \overset{(d)}{—} \boxed{\text{int. sol KdV}(s)} \overset{(e)}{—} \boxed{\text{Dym}(\rho)}$$

where all the third order nonlinear evolution equations are, respectively

$$\begin{aligned}
&u_t = u_{xxx} + 6uu_x && \text{(KdV)},\\
&v_t = v_{xxx} - 6v^2 v_x && \text{(mKdV)},\\
&w_t = w_{xxx} - 3\frac{w_x w_{xx}}{w} && \text{(new eq)},\\
&\varphi_t = \varphi_x\{\varphi; x\}, \quad \text{where} \quad \{\varphi; x\} := \left(\frac{\varphi_{xx}}{\varphi_x}\right)_x - \frac{1}{2}\left(\frac{\varphi_{xx}}{\varphi_x}\right)^2 && \text{(KdV sing.)},\\
&s^2 s_t = s^2 s_{xxx} - 3ss_x s_{xx} + \frac{3}{2}{s_x}^3 && \text{(int. sol KdV)},\\
&\rho_t = {\rho}^3 \rho_{\xi\xi\xi} && \text{(Dym)}.
\end{aligned}$$

The Bäcklund transformations linking these equations them, are, following their order in the Bäcklund chart:

$$(a)\ \ u + v_x + v^2 = 0, \qquad (b)\ \ v - \frac{w_x}{w} = 0, \tag{5.2}$$

$$(c)\ w^2 - \varphi_x = 0, \qquad (d)\ \ s - \varphi_x = 0, \tag{5.3}$$

and

$$(e)\ \ \bar{x} := D^{-1}s(x),\ \rho(\bar{x}) := s(x), \qquad \text{where} \qquad D^{-1} := \int_{-\infty}^{x} d\xi, \tag{5.4}$$

so that $\bar{x} = \bar{x}(s, x)$ and, hence, $\rho(\bar{x}) := \rho(\bar{x}(s, x))$. The transformation (e) denotes the reciprocal transformation, as it is termed to stress it interchanges the role of the dependent and independent variables[7] It represents a Bäcklund transformation between the extended manifold consisting of the both the dependent and the independent variables; namely, the manifold given by

[6] Combination of (1.12) with (1.19), respectively, Cole-Hopf and introduction of a *bona fide* potential when connecting the interacting soliton KdV with the KdV sing. equation [22] produce the transformation (5.1).

[7] see, for instance, [42] for a general introduction and application of reciprocal transformations. Details on the transformation (e) are given in [4, 22].

pairs formed by the dependent and the independent variables. Then, the transformation (e) defines, at least locally [4, 18] a Bäcklund transformation:

$$T_{(s,x)} \to T_{(\rho,\bar{x})}$$

between the two respective tangent spaces. Therefore, it is possible to transfer the infinitesimal structure using the transformation formulae [22, 18].

Now, taking into account the invariance under the Möbius group of transformations enjoyed by the singularity manifold equation, the Bäcklund chart can be duplicated to obtain

$$\begin{array}{ccccccccccc}
\boxed{\text{KdV}(u)} & \overset{(a)}{-} & \boxed{\text{mKdV}(v)} & \overset{(b)}{-} & \boxed{\text{new eq}(w)} & \overset{(c)}{-} & \boxed{\text{KdV sing.}(\varphi)} & \overset{(d)}{-} & \boxed{\text{int. sol KdV}(s)} & \overset{(e)}{-} & \boxed{\text{Dym}(\rho)} \\
AB_1 \updownarrow & & AB_2 \updownarrow & & AB_3 \updownarrow & & M \updownarrow & & AB_4 \updownarrow & & AB_5 \updownarrow \\
\boxed{\text{KdV}(u)} & \overset{(a)}{-} & \boxed{\text{mKdV}(v)} & \overset{(b)}{-} & \boxed{\text{new eq}(w)} & \overset{(c)}{-} & \boxed{\text{KdV sing.}(\varphi)} & \overset{(d)}{-} & \boxed{\text{int. sol KdV}(s)} & \overset{(e)}{-} & \boxed{\text{Dym}(\rho)}
\end{array}$$

where the vertical lines represent auto-Bäcklund transformations which are all induced by the combination of the Bäcklund transformations linking the other equations with invariance enjoyed by the KdV singularity equation. In detail, starting from the left hand side, the invariance AB_1 and AB_2 are the well known KdV and mKdV [35], auto-Bäcklund transformations, given in [2, 22], $\text{AB}_3 \equiv \text{I}$ is the invariance admitted by the novel nonlinear evolution equation (3.1) proved in Proposition 2. The last two, AB_4, and AB_5, respectively, are auto-Bäcklund transformations, [22], admitted by the int. sol. KdV and Dym equations. Notably, the connection between KdV and Dym equation [5, 22] finds applications in the construction of solutions admitted by the Dym equation [24, 26]. The extension to $2+1$ dimensional equations is given in [6, 36].

6 Admitted recursion operator & hierarchy

This section is devoted to the construction of the recursion operator admitted by the equation (3.1). Specifically, the Cole-Hopf link (3.5) between equation (3.1) and the mKdV equation allows to prove, according to [19, 18], it admits a recursion operator.

Proposition 3
The nonlinear evolution equation (3.1) *admis the recursion operator*

$$\Psi(w) = \frac{1}{2w} D w^2 \left[D^2 + 2U + D^{-1} 2 U D \right] \frac{1}{w^2} D^{-1} 2w \quad , \tag{6.1}$$

where

$$U := \frac{w_{xx}}{w} - 2\frac{w_x^2}{w^2} \quad . \tag{6.2}$$

Proof
Consider the Bäcklund transformation (3.5), then, the transformation operator Π, recalling (2.10), reads

$$\Pi := -B_w^{-1} B_\varphi \tag{6.3}$$

where

$$B_w[q] = \frac{\partial}{\partial\varepsilon}B(w+\varepsilon q,\varphi)\Big|_{\varepsilon=0} = \frac{\partial}{\partial\varepsilon}\left[(w+\varepsilon q)^2-\varphi_x\right]\Big|_{\varepsilon=0} = 2wq \qquad (6.4)$$

and,

$$B_\varphi[q] = \frac{\partial}{\partial\varepsilon}B(w,\varphi+\varepsilon q)\Big|_{\varepsilon=0} = \frac{\partial}{\partial\varepsilon}\left[w^2+(\varphi+\varepsilon q)_x\right]\Big|_{\varepsilon=0} = q_x \qquad (6.5)$$

hence

$$B_w = 2w \quad , \; B_\varphi = D \;\Rightarrow\; \Pi = -{B_w}^{-1}B_\varphi = \frac{1}{2w}D\ . \qquad (6.6)$$

Now, substitution of the latter in (2.9) gives

$$\Psi(w) = {B_w}^{-1}B_\varphi\Phi(\varphi)B_\varphi^{-1}B_w\big|_{w^2-\varphi_x=0}\,, \qquad (6.7)$$

where $\Phi(\varphi)$, according to formulae (1.23)-(1.24) in [22], is

$$\Phi(\varphi) = \varphi_x\left[D^2+\{\varphi;x\}+D^{-1}\{\varphi;x\}\right]\frac{1}{\varphi_x}D^{-1} \quad . \qquad (6.8)$$

Then, substitution of the latter and of the transformation operator Π in (6.7) gives the operator

$$\Psi(w) = \frac{1}{2w}D\varphi_x\left[D^2+\{\varphi;x\}+D^{-1}\{\varphi;x\}D\right]\frac{1}{\varphi_x}D^{-1}2w\Big|_{\varphi_x=w^2} \quad . \qquad (6.9)$$

recalling the definition $(3.4)_2$ of the Schwarzian derivative $\{\varphi;x\}$, on substitution of $\varphi_x = w^2$, it follows

$$\{\varphi;x\}|_{\varphi_x=w^2} = 2\left(\frac{w_{xx}}{w}-2\frac{w_x^2}{w^2}\right), \qquad (6.10)$$

the latter, on use of U, introduced in (6.2) to simplify the notation, gives

$$\{\varphi;x\}|_{\varphi_x=w^2} = 2U$$

and, hence,

$$\Psi(w) = \frac{1}{2w}Dw^2\left[D^2+2U+D^{-1}2UD\right]\frac{1}{w^2}D^{-1}2w \quad ,$$

which coincides with (6.1) and completes the proof. □

In addition, the following proposition holds.

Proposition 5

The recursion operator (6.1) *admitted by the nonlinear evolution equation* (3.1) *is hereditary.*

Proof
To prove the thesis, note that equation (3.1) is linked via Bäcklund transformations to all the nonlinear evolution equations in the Bäcklund chart; hence, since all of them admit a hereditary recursion operator, according to [18, 19], also the recursion operator (6.1) admitted by the newly obtained equation (3.1) enjoys the hereditariness property. □

Remark
The recursion operator (6.1) admitted by the equation (3.1) can be also obtained from the Cole-Hopf link $vw - w_x = 0$, with the mKdV equation, via the same method. Hence, in this case

$$\Psi(w) = {B_w}^{-1} B_v \Phi_{\mathrm{mKdV}}(v) B_v^{-1} B_w\big|_{vw-w_x=0}\,, \tag{6.11}$$

where, respectively, B_w and B_v are given by

$$B_w[q] = \frac{\partial}{\partial\varepsilon} B(w+\varepsilon q, \varphi)\Big|_{\varepsilon=0} = \frac{\partial}{\partial\varepsilon}\left[(w+\varepsilon q)^2 - \varphi_x\right]\Big|_{\varepsilon=0} = 2wq \tag{6.12}$$

and,

$$B_v[q] = \frac{\partial}{\partial\varepsilon} B(w, v+\varepsilon q)\Big|_{\varepsilon=0} = \frac{\partial}{\partial\varepsilon}\left[w(v+\varepsilon q) - w_x\right]\Big|_{\varepsilon=0} = wq \tag{6.13}$$

and

$$B_w[q] = \frac{\partial}{\partial\varepsilon} B(w+\varepsilon q, v)\Big|_{\varepsilon=0} = \frac{\partial}{\partial\varepsilon}\left[(w+\varepsilon q)v - (w+\varepsilon q)_x\right]\Big|_{\varepsilon=0} = vq - q_x \tag{6.14}$$

Explicit computations allow to obtain, once again, (6.1).

Now, equation (3.1) , when the hereditary recursion operator $\Psi(w)$ is given in (6.1), can be written as

$$w_t = \Psi(w) w_x \tag{6.15}$$

and the corresponding hierarchy is generated

$$w_t = [\Psi(w)]^n w_x \quad , \; n \in \mathbb{N}. \tag{6.16}$$

Since, as in [22], all the nonlinear evolution equations in the Bäcklund chart in Section 5 admit a hereditary recursion operator [18, 19], all the links can be extended to the corresponding whole hierarchies. Then, fixed $n = n_0, n_0 \in \mathbb{N}$ a different Bäcklund chart is obtained which links nonlinear evolution equations of order $2n_0+1$; the case $n_0 = 1$ corresponds to the 3rd order KdV-type equations; if $n_0 = 2, 3$, respectively, the nonlinear evolution equations in the Bäcklund chart are all of the 5th and 7th order.

The links among the corresponding members in the hierarchies can be depicted via the same Bäcklund chart in Section 5, that is, for each $n \in \mathbb{N}$, it holds

$$\boxed{[\Phi_1(u)]^n u_x} \overset{(a)}{-} \boxed{[\Phi_2(v)]^n v_x} \overset{(b)}{-} \boxed{[\Phi_3(w)]^n w_x} \overset{(c)}{-} \boxed{[\Phi_4(\varphi)]^n \varphi_x} \overset{(d)}{-} \boxed{[\Phi_5(s)]^n s_x} \overset{(e)}{-} \boxed{[\Phi_6(\rho)]^{n-1} \rho^3 \rho_{xxx}}$$

where, the recursion operators[8] are, respectively

$$\begin{aligned}
\Phi_1(u) \equiv \quad & \Phi_{\mathrm{KdV}}(u) = D^2 + 2DuD^{-1} + 2u && \text{(KdV)},\\
\Phi_2(v) \equiv \quad & \Phi_{\mathrm{mKdV}}(v) = D^2 - 4DvD^{-1}vD && \text{(mKdV)},\\
\Phi_3(w) \equiv \quad & \Psi(w) = \frac{1}{2w} Dw^2 \big[D^2 + 2U + D^{-1}2UD\big] \frac{1}{w^2} D^{-1} 2w && \text{(new eq)},\\
\Phi_4(\varphi) \equiv & \Phi_{\mathrm{KdVsing}}(\varphi) = \varphi_x \left[D^2 + \{\varphi; x\} + D^{-1}\{\varphi; x\}D\right] \frac{1}{\varphi_x} && \text{(KdV sing.)},\\
\Phi_5(s) \equiv \quad & \Phi_{\mathrm{KdVsol}}(s) = Ds \left[D^2 + S + D^{-1}SD\right] \frac{1}{s} D^{-1} && \text{(int. sol KdV)},\\
\Phi_6(\rho) \equiv \quad & \Phi_{Dym}(\rho) = \rho^3 D^3 \rho D^{-1} \rho^{-2} && \text{(Dym)},
\end{aligned}$$

where

$$U := \frac{w_{xx}}{w} - 2\frac{w_x^2}{w^2} \quad , \quad S := (\frac{s_x}{s})_x - \frac{1}{2}(\frac{s_x}{s})^2 \quad . \tag{6.17}$$

and the links among such hierarchies of nonlinear evolution equations, are indicated in (5.2).

Remark
The Dym hierarchy, generated on application of the hereditary recursion operator [2, 22, 31, 30] to the Dym equation,

$$\rho_t = [\Phi_{Dym}(\rho)]^n \rho^3 \rho_{xxx} \ , n \geqslant 0, \quad \text{where} \quad \Phi_{Dym}(\rho) = \rho^3 D^3 \rho D^{-1} \rho^{-2} \tag{6.18}$$

is connected to all the hierarchies in which appears in the Bäcklund charts in Section 5; hence, it is also related to the hierarchy of nonlinear evolution equation whose base member is (3.1).

7 Remarks, perspectives and open problems

This Section is devoted to collect some remarks and open problems which arise from the presented results. This study is stricktly connected to the research program which involves C.Schiebold and, lately also M.Lo Schiavo, and the author concerning operator evolution equations and their properties. The results already obtained, based on the theoretical foundation in [15] and references therein, concern KdV-type non-Abelian equations in [7, 8, 11, 13] where, analogies as well as some interesting differences which arise in

[8] The recursion operators here listed are well known ones with the only excepition of (6.1), see, for instance [2].

the non-Abelian case are poited out. Non-Abelian Burgers hierarchies are studied in [9, 10, 12]. Comments concerning the comparison non-Abelian vs. Abelian results are comprised in [7, 8, 3, 11, 12]. The only remarkable property to strees in the present contest is that, also in the non-Abelian case, hereditariness is preserved under Bäcklund transformations. Hence, proved the hereditariness of one recursion operator [47, 10], the hereditariness of all the recursion operator of other non-Abelian equations linked to it follows, see[44, 45, 46, 8, 11].

- The Bäcklund chart in Section 5 extends that one in [22, 37] finds its analogous Bäcklund chart in [4, 40], where 5th order nonlinear evolution equations, i.e. Caudrey-Dodd-Gibbon-Sawata-Kotera (CDG-SK) [14, 43] and Kaup-Kupershmidt (KK) equations, which, in turn, play the role of the KdV and mKdV equations appear. The 5th order nonlinear evolution equation analogous to the Dym eq is the Kawamoto equation [29]: it is linked via the reciprocal transformation (5.4) to the singularity manifold equation [49] related to the CDG-SK equation. However, further to the many analogies the transformations the two different Bäcklund charts are not exactly the same. Hence, the question arises whether or not there exist a 5th order analog of equation (3.1).
- All the structural properties which are preserved under Bäcklund transformations are enjoyed by all nonlinear evolution equations in the same Bäcklund chart. This is the case, in particular, of hereditariness of recursion operators. Also the Hamiltonian and/or bi-Hamiltonian structure [21, 23, 25, 32] in preserved under Bäcklund transformations. Hence, equation (3.1) is expected to admit a Hamiltonian and bi-Hamiltonian structure: the study is currently in progress.
- In [36] the $(2 + 1)$ dimensional KP, mKP and Dym hierarchies are connected via Bäcklund transformations; then in [37] it is shown that, when suitable constraints are imposed, the Bäcklund chart in [22] is obtained. A further question which arises is whether a $2 + 1$ equation analog of (3.1) can be constructed.

Further questions concerning open problems and perspective investigations in the case of operator equations are referred to [11].

Acknowledgements

The financial support of G.N.F.M.-I.N.d.A.M., I.N.F.N. and Sapienza University of Rome, Italy are gratefully acknowledged.

References

[1] F. Calogero and A. Degasperis, A modified modified Korteweg-de Vries equation, Inverse Problems **1**, (1985), 57-66. 1985 Inverse Problems 1 006

[2] F. Calogero and A. Degasperis, Spectral Transform and Solitons I, *Studies in Mathematics and its Application*, Vol. 13, North Holland, Amsterdam, 1980.

[3] S Carillo, Nonlinear Evolution Equations: Bäcklund Transformations and Bäcklund charts, *Acta Applicandae Math.* **122**, n. 1 (2012), 93-106.

[4] S. Carillo and B. Fuchssteiner, The abundant symmetry structure of hierarchies of nonlinear equations obtained by reciprocal links, *J. Math. Phys.* **30** (1989), 1606-1613.

[5] C. Rogers and M.C. Nucci, On reciprocal Bäcklund transformations and the Korteweg- de Vries hierarchy, Physica Scripta **33** (1986), 289-292.

[6] C. Rogers, The Harry Dym equation in 2+1-dimensions: a reciprocal link with the Kadomtsev-Petviashvili equation, *Phys. Lett.* **120A** (1987), 15-18.

[7] S. Carillo and C. Schiebold, Non-commutative KdV and mKdV hierarchies via recursion methods, *J. Math. Phys.* **50** (2009), 073510.

[8] S. Carillo and C. Schiebold, Matrix Korteweg-de Vries and modified Korteweg-de Vries hierarchies: Non-commutative soliton solutions, *J. Math. Phys.* **52** (2011), 053507.

[9] S. Carillo and C. Schiebold, A non-commutative operator-hierarchy of Burgers equations and Bäcklund transformations, in *Applied and Industrial Mathematics in Italy III: Selected Contributions from the 9th SIMAI Conference*, E. De Bernardis, R. Spigler e V. Valente Ed.s, SERIES ON ADVANCES IN MATHEMATICS FOR APPLIED SCIENCES, Vol.82, World Scientific Pubbl., Singapore, 2010, 175 -185.

[10] S. Carillo and C. Schiebold, On the recursion operator for the non-commutative Burgers hierarchy, *J.N.M.P.***19**1 (2012), 1250003, 11 pp.

[11] S. Carillo, M. Lo Schiavo and C. Schiebold, Bäcklund Transformations and Non Abelian Nonlinear Evolution Equations: a novel Bäcklund chart, *SIGMA* 12 (2016), 087, 17 pages.

[12] S. Carillo, M. Lo Schiavo and C. Schiebold, Recursion Operators admitted by non-Abelian Burgers equations: Some Remarks, *MATCOM*, in press (2017), arXiv: 1606.07270v1,12 pages.

[13] S. Carillo, M. Lo Schiavo and C. Schiebold, (2017), preprint.

[14] P. J. Caudrey, R. K. Dodd and J. D. Gibbon, A New Hierarchy of Korteweg-de Vries Equations, *Proc.R.Soc. London*, **A 351** (1976), 407-422.

[15] B. Carl and C. Schiebold, Nonlinear equations in soliton physics and operator ideals, *Nonlinearity* **12** (1999), 333-364.

[16] J.D. Cole, On a quasilinear parabolic equation occuring in aerodynamics, *Quart.App. Math.* **92** (1951), 25-236.

[17] D.A. Depireux and J. Schiff, On UrKdV and UrKP, *Lett. Math. Phys.* **33** (1995), 99-111.

[18] A.S. Fokas and B. Fuchssteiner, Bäcklund transformation for hereditary symmetries, *Nonlin. Anal., Theory Methods Appl.* **5** no. 4 (1981), 423-432.

[19] B. Fuchssteiner, Application of hereditary symmetries to nonlinear evolution equations, *Nonlin. Anal., Theory Methods Appl.* **3** no. 6 (1979), 849-862.

[20] B. Fuchssteiner, Solitons in interaction, *Progr. Theor. Phys.***78** no. 5 (1987), 1022-1050.

[21] B. Fuchssteiner, The Lie algebra structure of Degenerate Hamiltonian and Bi-Hamiltonian systems, *Progr. Theor. Phys.*, **68** (1982), 1082-1104.

[22] B. Fuchssteiner and S. Carillo, Soliton structure versus singularity analysis: Third order completely integrable nonlinear equations in 1+1 dimensions, *Phys. A* **154** (1989), 467-510.

[23] B. Fuchssteiner and S. Carillo, The Action-Angle Transformation for Soliton Equations, *Physica A*, **166** (1990), 651-676. doi:10.1016/0378-4371(90)90078-7.

[24] B. Fuchssteiner, T. Schulze and S. Carillo, Explicit solutions for the Harry Dym equation, *J. Phys. A* **25** (1992), 223-230.

[25] B. Fuchssteiner and W. Oevel, The bi-Hamiltonian structure of some nonlinear fifth- and seventh-order differential equations and recursion formulas for their symmetries and conserved covariants, *J. Math. Phys* **23** no. 3 (1982), 358-363.

[26] B.Yu Guo and C. Rogers, On the Harry Dym Equation and its Solution, *Science in China* 32 (1989), 283-295.

[27] C. Gu, H. Hu and Z. Zhou, Darboux transformations in integrable systems, Springer, Dordrecht, 2005, ISBN 1-4020-3087-8.

[28] E. Hopf, The partial differential equation $u_t + uu_x = muu_{xx}$, *Comm. Pure Appl. Math.* **3** (1950), 201-230.

[29] S. Kawamoto, An exact Transformation from the Harry Dym Equation to the modified KdV Equation, *J.Phys.Soc. Japan* **54** (1985),2055-2056.

[30] M. Leo, R.A. Leo, G. Soliani, and L. Solombrino, On the isospectral-eigenvalue problem and the recursion operator of the Harry Dym equation. Lett. Nuovo Cimento (2) **38** no. 2, (1983), 45?51.

[31] S.Y. Lou, Symmetries and similarity reductions of the Dym equation. *Phys. Scripta* **54** (1996), no. 5, 428-435.

[32] F. Magri, A simple model of the integrable Hamiltonian equation, *J.Math.Phys.* **19** (1978), 1156-1162.

[33] A.V. Mikhailov, A.B. Shabat, and R.I. Yamilov, The symmetry approach to the classification of non-linear equations. Complete lists of integrable systems Russian Math. Surveys **42**, 4 (1987), 1-63

[34] A.V. Mikhailov, V.S. Novikov, and J.P. Wang, (2007), On Classification of Integrable Nonevolutionary Equations. Studies in Applied Mathematics, **118**, 419-457.

[35] R.M. Miura Ed. Bäcklund Transformations, I.S.T. method and their applications, Lecture Notes in Math. 515, Springer, 1976.

[36] W. Oevel and S. Carillo, Squared Eigenfunction Symmetries for Soliton Equations: Part I, *J. Math. Anal. and Appl.* **217** no.1 (1998) 161-178. doi:10.1006/jmaa.1997.5707.

[37] W. Oevel and S. Carillo, *Squared Eigenfunction Symmetries for Soliton Equations: Part II*, *J. Math. Anal. and Appl.* **217** no.1 (1998) 179-199. doi:10.1006/jmaa.1997.5708.

[38] P.J. Olver, Evolution equations possessing infinitely many symmetries, *J. Math. Phys.***18** (1977), 1212-1215.

[39] C. Rogers and W.F. Ames, *Nonlinear Boundary Value Problems in Science and Engineering* Academic Press, Boston, 1989.

[40] C. Rogers and S. Carillo, On reciprocal properties of the Caudrey-Dodd-Gibbon and Kaup-Kupershmidt hierarchies, *Physica Scripta* **36** (1987), 865-869.

[41] C. Rogers and W.K. Schief, *Bäcklund and Darboux Transformations: Geometry and Modern Applications in Soliton Theory*, Cambridge University Press, Cambridge, 2002.

[42] C. Rogers and W.F. Shadwick. Bäcklund Transformations and Their Applications, *Mathematics in Science and Engineering* Vol. 161, Academic Press, Inc., New York-London, 1982.

[43] A.K. Sawada and A.T. Kotera, A method for finding N-soliton solutions of the KdV and KdV-like equations, *J. Progr. Theor. Phys.* **51** (1974), 1355-1367.

[44] C. Schiebold, Cauchy-type determinants and integrable systems, *Linear Algebra and its Applications* **433** (2010), 447-475.

[45] C. Schiebold, Noncommutative AKNS systems and multisoliton solutions to the matrix sine-Gordon equation, *Discr. Cont. Dyn. Systems Suppl.* **2009** (2009), 678-690.

[46] C. Schiebold, The noncommutative AKNS system: projection to matrix systems, countable superposition and soliton-like solutions, *J. Phys. A* **43** (2010), 434030.

[47] C. Schiebold, Structural properties of the noncommutative KdV recursion operator, *J. Math. Phys.* **52** (2011), 113504.

[48] J.P. Wang, A List of 1 + 1 Dimensional Integrable Equations and Their Properties, *J. Nonlinear Math. Phys.* **9**1, (2002), 213-233.

[49] J. Weiss, On classes of lntegrable systems and the Painlevé property, *J. Math. Phys.* **25** (1984), 13-24.

[50] G. Wilson, On the quasi-hamiltonian formalism of the KdV equation, *Physics Lett. A* **132** (1988), 445-450.